




Remote Sensing of Land

Homepage: www.gathacognition.com/journal/gcj1
<http://dx.doi.org/10.21523/gcj1>



Original Research Paper

Assessment of Aeolian Desertification Near Vedavathi River Cannel in Central Part of Andhra Pradesh: Remote Sensing Approach



Pradeep Kumar Badapalli, Raghu Babu K.*, Rajasekhar M. , Ramachandra M.

Department of Geology, Yogi Vemana University, Kadapa-516003, Andhra Pradesh (India).

Abstract

Desertification is one of the foremost environmental and social-economic problems facing the world. Thus, the observation and early detection of desertification is primary objective for scientific and policy organizations, with geospatial techniques as similar as Remote Sensing (RS) and Geographical Information System (GIS) being a candidate choice for the enlargement of monitoring systems. The present study purposes to assess the desertification caused by Aeolian sand deposits near Vedavathi River in a semi-arid region of Anantapur district, aided by temporal data of satellite imageries of past 28 years (1990 to 2018), were collected and processed. The extent of desertified land for the years 1990, 2000, 2010 and, 2018 is assessed, respectively about 28.22 Km², 33.86 Km², 46.30 Km² and, 58.62 Km². For the past 28 years, the Aeolian sand desertification is increased into the alarming rates with the land extent of 30.4 Km².

Article history

Received: 11 November, 2019;
 Final form: 06 February 2020;
 Accepted: 02 April 2020.

Keywords

Aeolian Sands;
 Desertification;
 GIS;
 Satellite Data;
 Semi-Arid Region.

Editor(s)

Suresh Kumar

© 2019 GATHA COGNITION® All rights reserved.

1 INTRODUCTION

Dry land of the world can be either classified as arid, established on climate or desert, established on surface physical characteristics (vegetation, landforms, etc.). For the persistence of this review, we will use the climatic classification implemented by UNESCO in 1979. Deserts thus can be categorized as semiarid having precipitation less than 550mm, arid with the precipitation less than 250mm and finally hyper arid with precipitation less than 25mm. Desertification is a foremost type of land deprivation happening around the globe, has convert one of the thoughtful conservational issue to the world (Reynolds *et al.*, 2007; Adger *et al.*, 2001; Tucker *et al.*, 1991). The UNCOD [United Nations Conference on Desertification] conference held at Nairobi in the year 1977 pronounced biological impending reasons for desertification. Afterwards, the fruition of the desertification and its definition have been reviewed by many scholars (Pradeep Kumar *et al.*, 2019; Amal Kar, 2016; Arya *et al.*, 2014; Bai *et al.*, 2011; Ajai *et al.*, 2009; Reynolds *et al.*, 2007;

Anonymous, 2005; Reynolds and Stafford-Smith, 2002; Eswaran *et al.*, 2001; Thomas and Meddleton, 1994; Rajasekhar *et al.*, 2020). However greatest inclusive and extensively acknowledged explanation of 'desertification' denotes to land deprivation is happening in arid, semi-arid, and sub-humid zones, consequential after the influences with climatic variations then humanoid doings (UNEP, 1992). The Aeolian desertification is a foremost kind of desertification; it is well-defined as land degradation by wind erosion, mostly subsequent from extreme humanoid doings and climate variation in deserted regions (UNCCD, 2004). At universal and provincial regions, wind plays significant part in geomorphological progressions and environmental alteration due to erosion, transportation and deposition of superficial sediments by the exploit of wind is the key for Aeolian process. Consequently, after the standpoint of geomorphology, land degradation progression in which wind movement is the important forceful influence, and

* Author address for correspondence

Department of Geology, Yogi Vemana University, Kadapa-516003, Andhra Pradesh (India).

Tel.: +91 9440673734

Emails: badapallipradeep@gmail.com (Pradeep Kumar B.); dr.kraghu@gmail.com (Raghu Babu K. -Corresponding author); sekhar.raja042@gmail.com (Rajasekhar M.); ramachandramesa@gmail.com (Ramachandra M.).

<https://dx.doi.org/10.21523/gcj1.19030104>

© 2019 GATHA COGNITION® All rights reserved.

sandy desertification (Eswaran *et al.*, 2001, Rajasekhar *et al.*, 2019b). The study part of the Anantapur district is one of the most sensitive environments and is suffering wide-ranging draught environments and Aeolian desertification (Pradeep Kumar *et al.*, 2019; Rajasekhar *et al.*, 2019c). This area is strongly affected by regional atmospheric circulation patterns and having low rainfall of about 520mm considered as lowermost in the study area. Most regions of the present study grieve from an arid climate, this ensuing in a delicate ecology that generates advantageous circumstances for desertification (Mas *et al.*, 1999; Li *et al.*, 2001). Under the impact of climate change and human activates, this become significant (Harshika *et al.*, 2012). Desertification has become severe problem that is obstructing socio-economic expansion and threatening its ecological environments of the study part (Zhang *et al.*, 2003). Therefore, present study attempted to detect and delineate the desertified land near to Vedavathi (Hagari) River in Adrea Pradesh (India) using Landsat datasets.

2 STUDY AREA

The Vedavathi (Hagari) River is the tributary of the Tunga Bhadra River. The Vedavathi River rises from the Bababudanagiri Mountains of Western Ghats and

streams through the Karnataka and Andhra Pradesh states. The river Vedavathi known as Hagari in the lower reaches rises near in the Western Ghats. It has a total length of 391 Km and of which the length in the Karnataka state is 293 Km and forms the common boundary between Karnataka and Andhra Pradesh for 26 Km. Vedavathi river lies between 14°30' and 15°10' N and 76°45' and 77°15'E and mainly flows through Bommanahal, Kanekal, Beluguppa, Brahmasamudram, Gummagatta, Rayadurgam, D-Hirehal mandalas of Anantapur district which is the South most district of Rayalaseema region of Andhra Pradesh state and it is fallen in the Survey of India (SOI) toposheets of 57B/13, 57B/14, 57B/15, 57F/1, 57F/2, 57F/3, 57F/5, 57F/6 and 57F/7 (Figure 1). The study area west being located in the rain shadow area of Western Ghats and is one among the villages which are frequently facing drought conditions in the country. The present study area receives an average rainfall of 520mm is lower most in the state (Pradeep Kumar *et al.*, 2018, Rajasekhar *et al.*, 2018a and 2018b). This region is facing humid climate with temperature alternating from 24 to 46° C in summer. Vedavathi River is the foci and most significant rivers in Anantapur district. It is seasonal or an ephemeral river, thus remain as dry throughout the




Figure 1. Study area: Vedavathi River

year. Local historical evidence shows that, nearly 200 years ago, due to heavy flood in the Vedavathi River, maximum of the fine sand from the river bed has been drifted and placed up to a substantial distance on agricultural fields plummeting soil fertility on the eastern sideways of the river. During the months of June to August, this region experiences strong surface winds nearly 36 to 38 Km/h from west to easterly direction by the Southwest Monsoon. Unidirectional Asymmetrical Ripple marks in the sand dunes are evidence for this migration in easterly direction of the study part. Figure 5 depicts the Anemometer wind speed in the SW monsoon). For this durable surface winds sand has been wandered to the crop fields and the land degradation is happening and finally decertified.

2.1 Hydrogeomorphology Setting

The study area covered by hard rock terrain along with an Archaean Peninsular Gneissic Complex (PGC) of hornblende-biotite gneiss, granodiorite and Lamprophyre (GSI, 2002). These granite rocks intersect with the underground dikes. The major rock types

present in the eastern portion of the study area belongs to the Archean age, they are PGC consisting felsic hornblende-biotite gneiss, hornblende gneiss, granite, granodiorite, biotite gneiss and Migmatites. Quartzite, Banded Iron Formations (BIF), Banded Magnetite Quartzite's and ferruginous quartzite's are also present in the top portion of the study area, they belong to the Dharwar super group of Archean to lower Proterozoic age. Pink granite or Grey granite (closepet granite) is current in the western portion of the study area (Figure 2a). In case of denudational landforms, shallow weathered pediplain, fairly and shallow weathered pediplains, pediment, pediment complicated, residual hill and denudational hill, fairly and shallow weathered pediplains, pediment, pediment in complicated and active geomorphological changes have been occurred by the action of wind or Aeolian process. Unidirectional Asymmetrical Ripple marks are present in the study part, which gives evidence for the active Aeolian process (Figure 7). Unstable sand dunes and alluvium is seen all along the Vedavathi River (Figure 2b).




Figure 2a. Geology




Figure 2b. Geomorphology

3 METHODOLOGY

3.1 Database

The spatial database consisted of change detection layers prepared from RS data Landsat 4, 5, 7 and 8 satellite imageries are used. The satellite data is geometrically corrected enhanced for visual interpretation, ground truth collection and land degradation map preparation. Landsat 8 satellite imagery merged satellite data is available as a default base layer which is helpful in delineating the change classes in a better way (Table 1). Landsat 4-5, Landsat 7 satellite images for year 1990 to 2010 and Landsat 8 satellite images for 2018 were used for LULC change detection. These images were imported in ERDAS Imagine software 2014, ArcGIS environment to prepare FCC [False Color Composite] images. The sub-setting of data sets were executed for extracting present study from satellite images by implementing georeferencing for boundary outline of area of interest.

3.2 Image Pre-processing

LULC classification carried out through supervised image classification in ERDAS IMAGINE 2014 software, using Landsat satellite imageries over the study area, which were acquired from USGS Earth Explorer (<https://earthexplorer.usgs.gov/>). In the present study, an integrated use of multispectral satellite data is used for the generation of database and extraction of numerous LULC patterns. Details of the data used are shown in Table 1. The image classification process performed in ERDAS imagine environment to prepare land use / land cover (LULC) analysis by supervised classification. Four signature files have been created namely sand/sand dunes, vegetation, fallow land and water bodies (Table 3). These supervised classifications based LULU has been done for the years 1990, 2000, 2010 and, 2018. Vedavathi River is attentive, where the sand is initiated and migrated from the tributary banks to adjoining areas. Vedavathi river shape file is created and clipped on LULC map for the past 28 years. LULC parameters were analyzed and classified into 5 classes, associated with their orientation in space (Rajasekhar et al., 2019c). The determination of the Landsat images was impacted by low cloud cover over the study area (Table 1). These images are pre-processed by clip methods using geospatial techniques such as, ERDAS Imagine 2014 and ArcGIS environments.

3.3 Land Use Land Cover Classification

To carry out the LULC analysis, supervised classification technique with maximum likelihood

algorithm was applied in the ERDAS Imagine 2014. Maximum likelihood algorithm is one of the best and easiest approaches in the image classification method with RS data sets. This approach is based on the pixel based probability analysis for a specific feature. The theory adopts for probabilities are identical for all features and the input pixels have a standard dissemination. However, this method takes a long time to calculate, is highly dependent on the normal distribution of data in each input range, and tends to over classify with relatively signature editor in the covariance matrix. The spectral analysis approach computes the spectral analysis between the candidate vector pixel measurement vector and the mean vector for each signature, and the spectral analysis (Table 2). It requires the shortest calculation time among other image classification approach, but pixels that do not need to be unclassified are classified and class variability is not taken into account. Field work was carried out for inaccessible and doubtful regions were corrected in the ERDAS Imagine environment. The most important final step at classification process is accuracy assessment. The aim of accuracy assessment is to quantitatively assess how effectively the pixels were sampled into the correct land cover classes. Moreover the key emphasis for accuracy assessment pixel selection was on areas that could be clearly identified on Landsat image, Google earth images and Google map. Four LULC change detection types are classified in the study area viz., (i) sand dunes (ii) vegetation (iii) fallow land and (iv) water body.

3.4 Change Detection Analysis

Geometric calculation method is also adopted for knowing the area of desertification or migrated area. The collective data has been handled for preparation of Desertification Status Maps (DSM). Further, we have collected statistical data for perceptive the rate of relocation of Aeolian sands. For execution change detection, a supervised and unsupervised classification was performed in the ERDAS imagine software by image classification technique. A pixel-based judgment was used to prepared change detection by pixel data to interpret the changes more efficient for the advantage of 'from' 'to' data. The classified images for various years' data compared using pixel data calculation to identify to calculative aspects in years from 1990 to 2018 (Rajasekhar et al., 2018b). Computable areal data of the overall desertified changes as well as gains and losses in both class between 1990 and 2018 were then accumulated. The overall methodology for present study is shown Figure 3.




Figure 3. Methodology

Table 1. Datasets

Datasets	Data used	Spatial resolution	Years	Source
Satellite data	Landsat 4-5	30m	1990	http://earthexplorer.usgs.gov/
	Landsat 7	30m	2000	http://bhuvan.nrsc.gov.in
	Landsat 8	30m	2010	
	Landsat 8	30m	2018	
Secondary data	Topographic maps 57B/13, 57B/14, 57B/15, 57F/1, 57F/2, 57F/3, 57F/5, 57F/6, 57F/7 Meteorological data		2000-2019	Survey of India MOSDAC
Primary data	Climate, soils, etc.			Groundwater Department, Anantapur District, Andhra Pradesh
Equipment used	Anemometer used for measuring wind speeds			

4 RESULTS AND DISCUSSIONS

4.1 Land Use Land Cover

LULC in the region includes vegetation, fallow land, water bodies, and sand dunes. Present study shows LULC is major result to identify and estimate the land use exploitation which has a greater key role in the desertification by man-made process (Sreenivasulu *et al.*, 2014; Rajasekhar *et al.*, 2017 and 2019c; Pradeep Kumar *et al.*, 2019). They have also contributed to the reduction of biological productivity, diversity and sustainability of the land. Mining, brick kiln deeds and industrial seepage affected areas are included under the anthropogenic factors (Figure 4).

In this part an active geomorphological changes have been occurred by the action of wind. Unidirectional Asymmetrical Ripple marks are present in the study part, which gives evidence for the active desertification (Figure 7). Unstable sand dunes and alluvium is seen all sideways of the Vedavathi river due to high surface winds happening from SW monsoon (June -September). This sand has been migrated, the land degradation happens and finally desertified. Sand and dunes play an important key role in the change of ecosystems in the present study and its categorized into different types with their size, shape, incidence of environment, degree of complexity and the direction of winds. The risk of wind erosion is severe in the arid and semi-arid regions. It includes both the removal and deposition of soil particles due to the action of the wind and the abrasive effects of moving particles during transport. In sandy areas that accumulate with rain, the sand partially stabilizes completely, depending on the vegetation cover that it establishes.

4.2 Change Detection Analysis

The Vedavathi River areas are fallen under the rain shadow zone according to the Koeppen's climatic division in India. Migration of the desert sand dunes are usually seen in the right side of the mandalas. Vedavathi river bed is the place of origination of the sands and the migration is carried by the transporting agent of wind. Hence these sands are called as Aeolian sands and this type of desertification is named as Aeolian desertification. The south-west monsoon season (June-September), study area facing high surface winds flow West to Eastern side, with the speed of 22 Km/h to 36 Km/h (Figure 5). For of these strong surface winds, the sand has been migrated to the agricultural fields which is reducing the soil fertility and causing devastation to the agriculture in the study area. The resultant figure 5 clearly depicts the rate of migration of sand and sand dunes in the study area. Different size and shapes of the

sand dunes present in the study part, wherever transverse and linear type of dunes are very common. Figure 7 shows different types of sand dunes and desertification status in the study area. Unidirectional Asymmetrical Ripple marks will give the evidence of this migration of sands and desertification from west to eastern side. All approaches for the past 28 years have been taken in to the consideration, i.e., 1990 to 2018, the analyzed statistics clearly reveals the rate of migration of Aeolian sands along the Vedavathi River. Present study indicates the DSM individually for the years 1990, 2000, 2010, and 2018 (Figure 6). In the year 1990 Aeolian sand desertification is calculated in the extent of 28.22 Km² and in the year 2000 it is calculated in the extent of 33.86 Km². For the decade 1990 to 2000 desertification is increased to 5.64 Km². In the year 2010 the Aeolian sand desertification is calculated in the extent of 46.30 Km² and in the year 2018 it is calculated in the extent of 58.62 Km² (5862 hectares). For the year 2010 to 2018 the desertification is increased to 12.32 Km². For the past 28 years the Aeolian sand desertification is increased in to the alarming rates with the land extent of 30.4 Km² (Table 3). Migration of sand in 28 years' span is calculated through geometric calculation technique, which evidently expose that the sand dunes were migrated uninterruptedly and desertification process is highly active in this area. This study clearly reveal that the Aeolian sands present in the Vedavathi River is caused for this desertification.



4.3 Accuracy Assessment

In grouping procedure, the role of Accuracy Assessment (AA) is quite vital. For the significant accentuation for exactness evaluation, pixel choice was on zones that could be obviously distinguished on Landsat high resolution image, Google Earth and Google Map (Schull *et al.*, 2005). Image was characterized at 50 points. Topographical guides and Google Earth maps were used as a kind of perspective source to group the selected features (Dewan and Yamaguchi, 2016). Table 2 demonstrates the connection between ground truth information and the relating characterized information, revealed through the error matrix report. The accuracy can be computed by:

$$\text{Overall accuracy} = \frac{\text{No. of correct points}}{\text{Total number of points}} \% 100$$

KAPPA analysis is a discrete multivariate technique used in accuracy assessments (Jenness and Wynne, 2007). KAPPA analysis yields a Khat statistic (an estimate of KAPPA) that is a measure of agreement or accuracy. Using the formulae equipped on ArcGIS. The results from the AA showed overall accuracy about 84% from random sampling process for the image.

Table 2. Ground signatures of Landsat imageries

Category	Landscape	Tone and texture	Picture of landscape	FCC subset
Sand/sand dunes	Sand dunes, including areas with flat sands, dense barchans, dues chains, mobile star dunes, and climbing dunes	Whitish yellow to bright white and wavy texture.		
Vegetation	The form and branching of the plants and trees and scrub will determine.	Vegetation appears in red color, this is more suitable in vegetation identification.		
Fallow land	Agricultural fields without crops or sometimes waste lands may include as fallow land.	Bluish/greenish grey with smooth texture. Lighter tone with irregular texture		
Water body	It comprises man-made reservoirs / lakes / tanks / canals also natural lakes, rivers / streams and creeks, etc.	Cyanish blue to blue according to the depth of water and sediment content with dark tone.		




Figure 4. Land use land cover

Table 3 Change detection analysis

Years	1990	2000	2010	2018	Change (1990-2018)
Desertified area (Km ²)	28.22	33.86	46.30	58.62	30.4




Figure 5. Wind speeds




Figure 6. Change detection




Figure 7. **(a)** Anemometer readings; **b, c, d and e:** Asymmetrical ripple marks; **f, g, h, and i;** Sand dunes encroachment in the crop land

5 CONCLUSIONS

Landsat data from 1990 to 2018 has been castoff to examine the movements in the desertification sensitivity over the earlier twenty-eight years along the Vedavathi River. This work discloses that migration of sand dunes and desertification happening by the Aeolian sand deposits present in the Vedavathi River. RS Landsat data has been effectively helped to develop DSM maps. The current research exposes that Aeolian desertification in the present study growing extremely fast by the transformation in environmental conditions. In the past 28 years 30.4 km² of agricultural land is desertified. Bommanahal, Kanekal, and Beluguppa mandalas are seriously affected by the Aeolian desertification, which are present in the eastern side of the Vedavathi River in south central part of the Anantapur district of AP, India. For monitoring the desertification and to take action plans towards the mitigation of desertification is possible by DSM maps. Some of the recommendations can comprise that to decrease of surface wind speeds, in order to sidestep sand migration on agricultural fields, suitable agriculture aid to inhibit the desertification and restoration of rigorously desertified lands for prolific consumption for cultivation perseverance.

ACKNOWLEDGEMENTS

The first author B. Pradeep Kumar, greatly thankful to Department of Science and Technology (DST), Government of India, for financial support through INSPIRE programme (Sanction order No. DST/INSPIRE Fellowship/2017/IF170114). We are also thankful to USGS for remote sensing data and Department of Geology, Yogi Vemana University, for necessary facilities for carrying out the research work.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ABBREVIATIONS

AA: Accuracy Assessment; **BIF:** Banded Iron Formations; **DSM:** Desertification Status Maps; **FCC:** False Colour Composite; **LULC:** Land Use Land Cover; **GIS:** Geographical Information System; **GSI:** Geological Survey of India; **PGC:** Peninsular Gneissic Complex; **RS:** Remote Sensing; **SOI:** Survey of India; **UNCOD:** United Nations Conference on Desertification.

REFERENCES

Adger, W. N., 2001. Scales of governance and environmental justice for adaptation and mitigation of climate change. *Journal of International Development*, 13(7), 921-931. DOI: <https://doi.org/10.1002/jid.833>

Ajai, Arya, A., Dhinwa, P., Pathan, S., and Raj, K., 2009. Desertification/Land degradation status mapping of India. *Current Science*, 97, 25.

Alkharabsheh, M. M., Alexandridis, T. K., Bilas, G., Misopolinos, N. and Silleos, N., 2013. Impact of land covers change on soil erosion hazard in northern Jordan using remotesensing and GIS. *Procedia Environmental*

Sciences, 19, 912-921. DOI: <https://doi.org/10.1016/j.proenv.2013.06.101>

Amal, K., 2016. Desertification, Central Arid Zone Research Institute (CAZRI).

Anonymous, 2003. Desertification Status Mapping- Technical Guidelines. Forest, Land use and Photogrammetry group, RESIPA, Space Application Venter, Ahmadabad, 34.

Arya, V. S., Hardev Singh, Hooda, R. S. and Arya, A. S., 2014. Desertification change analysis in Siwalik hills of Haryana using Geo-Informatics. The international archives of the Photogrammetry, Remote Sensing And Spatial Information Sciences, XL-8. DOI: <https://doi.org/10.5194/isprsarchives-XL-8-677-2014>

Bai, Z., Dent, D., Wu, Y. and de Jong, R., 2013. Land degradation and ecosystem services. In *Ecosystem services and carbon sequestration in the biosphere*, 357-381. Springer, Dordrecht. DOI: https://doi.org/10.1007/978-94-007-6455-2_15

Budihal, S. L., Raj, K. G., Reddy, R. S., Natarajan, A., Paul, M. A. and Bandyopadhyay, S., 2006. Assessment and mapping of desertification status in Bellary district, Karnataka State, using IRS data. In *ISPRS Commission IV International Symposium on Geospatial Databases for Sustainable Development*, Goa, 25-30.

Dhinwa, P. S., Dasgupta, A. and Ajai, 2016. Monitoring and assessment of desertification using satellite remote sensing. *Journal of Geomatics*, 10(2), 201-216.

Eswaran, H., Lal, R. and Reich, P. F., 2001. Land degradation: An overview. *Responses to Land degradation*, 20-35.

GSI [Geological Survey of India], 2002. Geological Survey of India District Mineral Resources Map Anantapur, Andhra Pradesh, India.

Harshika, A. K. and Sopan I., 2012. Land use land cover classification and change detection using high resolution temporal satellite data. *Journal of Environment*, 01(04), 146-15.

Jenness, J. and Wynne, J. J., 2007. Kappa analysis (kappa_stats. avx) extension for ArcView 3x. Jenness Enterprises.

Lam, D. K., Rimmel, T. K., Drener, T. D., 2011. Tracking desertification in california using remote sensing: A sand dune encroachment approach. *Remote sens*, 3, 1-13. DOI: <https://doi.org/10.3390/rs3010001>

Li, X. J., Li, Q. S., Wang, Z. C., Liu, X. T., 2002. A research on characteristics and rational exploitation of soda saline land in the Western Songnen Plain. *Research of Agricultural Modernization*, 23, 361-364.

Mas, J. F., 1999. Monitoring land-cover changes: a comparison of change detection techniques. *International Journal of Remote Sensing*, 20(1), 139-152. DOI: <https://doi.org/10.1080/014311699213659>

Pradeep Kumar, B., Raghu Babu, K., Rajasekhar, M. and Ramachandra, M., 2019. Assessment of land degradation and desertification due to migration of sand and sand dunes in Beluguppa Mandal of Anantapur district (AP, India), using remote sensing and GIS techniques. *The Journal of Indian Geophysical Union*, 23(2), 173-180.

Pradeep Kumar, B., Rajasekhar, M., Ramachandra, M. and Siva Kumar Reddy P., 2018. Assessment of land degradation and desertification due to migration of sand dunes - A case study in Bommanahal Mandal, Anantapur district, Andhra Pradesh, India using Remote Sensing and GIS techniques. *IJRAT*, 6 (6), 1-7.

Pradeep Kumar. B., Raghu Babu. K., Rajasekhar M. and Narayana Swamy, B., 2019. Change detection analysis of Land Use and Land Cover in Tadimarri mandal of

- Anantapur district, AP, India, using Geospatial Techniques. *IJRAR- International Journal of Research and Analytical Reviews*, 6(2), 634-640.
- Raghu Babu, K. and Sudharshan Raju, G., 2012. Application of Remote Sensing for delineation of Uranium bearing Vempalli dolomites in and around Tummalapalli area, Cuddapah Basin, India. *International Journal of Geomatics and Geosciences*, 2(3), 842-852.
- Rajasekhar, M., Sudarsana. Raju, G., Siddi Raju, R., Ramachandra, M. and Pradeep Kumar, B., 2018b. Data in brief data on comparative studies of lineaments extraction from ASTER DEM, SRTM and Cartosat for Jilledubanderu River basin, Anantapur district, A. P, India by using remote sensing and GIS. *Data in Brief*, 20, 1676-1682. DOI: <https://doi.org/10.1016/j.dib.2018.09.023>
- Rajasekhar, M., Gadhiraaju, S. R., Kadam, A. and Bhagat, V., 2020. Identification of groundwater recharge-based potential rainwater harvesting sites for sustainable development of a semi-arid region of southern India using geospatial, AHP, and SCS-CN approach. *Arabian Journal of Geosciences*, 13(2), 24. DOI: <https://doi.org/10.1007/s12517-019-4996-6>
- Rajasekhar, M., Raju, G. S. and Raju, R. S., 2019a. Assessment of groundwater potential zones in parts of the semi-arid region of Anantapur District, Andhra Pradesh, India using GIS and AHP approach. *Modeling Earth Systems and Environment*, 5(4), 1303-1317. DOI: <https://doi.org/10.1007/s40808-019-00657-0>
- Rajasekhar, M., Raju, G. S., Raju, R. S. and Basha, U. I., 2018a. Data on artificial recharge sites identified by geospatial tools in semi-arid region of Anantapur District, Andhra Pradesh, India. *Data in brief*, 19, 462-474. DOI: <https://doi.org/10.1016/j.dib.2018.04.050>
- Rajasekhar, M., Raju, G. S., Raju, R. S. and Basha, U. I., 2017. Landuse and landcover analysis using remote sensing and GIS: A case study in Uravakonda, Anantapur District, Andhra Pradesh, India. *IRJET*, 4(9), 780-785.
- Rajasekhar, M., Raju, G. S., Sreenivasulu, Y. and Raju, R. S. 2019b. Delineation of groundwater potential zones in semi-arid region of Jilledubanderu river basin, Anantapur District, Andhra Pradesh, India using fuzzy logic, AHP and integrated fuzzy-AHP approaches. *HydroResearch*, 2, 97-108. DOI: <https://doi.org/10.1016/j.hydres.2019.11.006>
- Rajasekhar, M., Sudarsana Raju, G., Bramaiah, C., Deepthi, P., Amaravathi, Y. and Siddi Raju, R., 2018c. Delineation of groundwater potential zones of semi-arid region of YSR Kadapa District, Andhra Pradesh, India using RS, GIS and analytic hierarchy process. *Remote Sensing of Land*, 2(2), 76-86. DOI: <https://doi.org/10.21523/gcj.1.18020201>
- Rajasekhar, M., Sudarsana Raju, G., Imran Basha, U., Siddi Raju, R., Pradeep Kumar, B., and Ramachandra, M. 2019c. Identification of suitable sites for artificial groundwater recharge structures in semi-arid region of Anantapur district: AHP approach. *Hydrospatial Analysis*, 3(1), 1-11. DOI: <https://doi.org/10.21523/gcj.3.19030101>
- Reynolds, J. F. and Stafford Smith, D. M., 2002. Do humans cause deserts? An old problem through the lens of a new framework: The dahlem desertification paradigm, *Proceedings of the VIIth International Rangelands Congress*, 2042-2048.
- Reynolds, J. F., Maestre, F. T., Kemp, P. R., Stafford-Smith, D. M. and Lambin E., 2007. Natural and human dimensions of land degradation in drylands: Causes and consequences. In: Canadell J. G., Pataki D.E., Pitelka L.F. (eds) *Terrestrial Ecosystems in a Changing World*. Global Change - The IGBP Series. Springer, Berlin, Heidelberg.
- Schull, M. and Turral, H., 2005. Ganges and Indus river basin land use / land cover (LULC) and irrigated area mapping using continuous streams of MODIS data, 95, 317-341. DOI: <https://doi.org/10.1016/j.rse.2004.12.018>
- Sreenivasulu, G., Jayaraju N. and Lakshmi P. T., 2014. Landuse and landcover analysis using remote Sensing and GIS: A case study in and around Rajampet, Kadapa District, Andhra Pradesh, India. *IJMRD*, 2(1), 186-191.
- Thomas, D. S. G. and Middleton, N. J., 1994. *Desertification: Exploding the Myth*. John Wiley & Sons Ltd., Chichester, UK.
- Tucker, C. J., Dregne, H. E. and Newcomb, W. W., 1991. Expansion and contraction of the Sahara Desert from 1980 to 1990. *Science*, 253, 299-300. DOI: <https://doi.org/10.1126/science.253.5017.299>
- UNCCD [UN Conference on Desertification], 1978. Round up, plan of Action and Resolutions, New York.
- UNEP, 1992. Desertification, Land degradation [definitions]. Desertification Control Bulletin 21.
- Zhang, X. Y., Gong, S. L., Zhao, T. L., Arimoto, R., Wang, Y. Q. and Zhou, Z. J., 2003. Sources of Asian dust and role of climate change versus desertification in Asian dust emission. *Geophysical Research Letters*, 30(24), 8.1-8.4. DOI: <https://doi.org/10.1029/2003GL018206>
