



Hydrospatial Analysis

Homepage: www.gathacognition.com/journal/gcj3
<http://dx.doi.org/10.21523/gcj3>



Original Research Paper

Groundwater Quality Analysis of an Emerging Part of Suburb of Pune Metropolitan Region, Maharashtra, India using GIS and Remote Sensing Techniques



Satyajit K. Gaikwad^{1*}, Suryakant P. Gaikwad¹, Dnyaneshwar V. Date², Somnath T. Borhade², Avinash M. Kandekar², Ashwini H. Supekar¹, Ravindra S. Bhagat³, Pravin N. Kambale⁴, Ramdas B. Madale⁵, Tushar P. Raut⁶, Omkar S. Kadekar¹

1. Department of Geology, Savitribai Phule Pune University, Pune - 411 007 (India).
2. Department of Geography, Savitribai Phule Pune University, Pune - 411 007 (India).
3. Department of Geography, S. P. M. Sahebrao Shankarrao Dhamdhare Arts and Commerce College, Talegaon Dhamdhare, Pune, Maharashtra (India).
4. Department of Geography, Dr. Babasaheb Ambedkar Commerce and Maharshi Vitthal Ramji Shinde Arts College Nanapeth Pune, Maharashtra (India).
5. Department of Geography, Swami Vivekanand College, Mukrabad, Mukhed, Dist. Nanded, Maharashtra, India.
6. Department of Geography, C.T. Bora College, Shirur, District – Pune. Maharashtra, India.

Abstract

Thirty (30) groundwater samples have been collected during Pre-monsoon season-2015 to analyze the groundwater quality of Shikrapur and Talegaon Dhamdhare area. Maps were prepared for major physicochemical elements in groundwater and geomorphologic aspects using GIS and Remote Sensing techniques. Trend in cations is $\text{Na} > \text{Ca} > \text{Mg} > \text{K}$ while in anions is $\text{Cl} > \text{HCO}_3 > \text{NO}_3 > \text{SO}_4$. The average, $\text{Na} + \text{Ca}$ representing 61.37% of total cations denoting major supply from weathering of plagioclase feldspar while $\text{Ca} + \text{Mg}$ values, 67.92 % of cations contributed from olivine and pyroxene. Anions like Cl , SO_4 and NO_3 in groundwater is contributed from anthropogenic activities.

The results were compared with WHO norms and found higher values for Electrical Conductivity (EC), Calcium (Ca), Potassium (K), Magnesium (Mg), Chloride (Cl) and Nitrate (NO_3). Other elements show low to optimum values indicating good quality for drinking purposes, excluding some pockets from lower reaches. Regular quality monitoring for groundwater with rainwater harvesting is suggested to improve the quality of groundwater in the region.

Article history

Received: 12 January 2019
 Revised: 18 February 2019
 Accepted: 16 March 2019

Keywords

Groundwater Quality;
 Metropolitan Region;
 GIS;
 Remote Sensing;
 Pune City.

Editor(s)

P. S. Hire

© 2018 GATHA COGNITION® All rights reserved.

1 INTRODUCTION

Globally, groundwater is the major source for domestic, agriculture as well as for industrial purposes. (Kale et al., 2010; Ravishankar et al., 2010; Gaikwad et al., 2012a; Ravikumar and Somashekar, 2013; Wagh et al., 2016; Wagh et al., 2018a; Wagh et al., 2018b; Gaikwad et al., 2019). Major elements like sodium (Na^+), calcium (Ca^{2+}), magnesium (Mg^{2+}), potassium (K^+), bicarbonate (HCO_3), chloride (Cl^-), sulphate (SO_4^{2-}), nitrate (NO_3^-)

and silica (Si) makes up to 99% of the dissolved constituents in groundwater (MacDonald and Davies, 2000; Ganyaglo et al., 2010). Once groundwater aquifers contaminated by anthropogenic activities, it will persist that contamination for hundreds of years due to slowest movement (Nathanson, 1986; Aksoy and Scheytt, 2007; Ravishankar et al., 2010).

* Author address for correspondence

Department of Geology, Savitribai Phule Pune University, Pune - 411 007 (India).
 Tel.: +91 9881735343

E-mails: satyajit.gaikwad@gmail.com (S. Gaikwad -Corresponding author); suryakantgaikwad7@gmail.com (S. Gaikwad); dnyanandate@gmail.com (D Date); somashu.borhade7421@gmail.com (S. Borhade); amkandekar@unipune.ac.in (A. Kandekar); ashwinis@unipune.ac.in (A. Supekar); ravindra.bhagat1@gmail.com (R. Bhagat); praveenkamble@gmail.com (P. Kambale); rdmadale@gmail.com (R. Madale); tusharraut103@gmail.com (T. Raut); komkar.3005@gmail.com (O. Kadekar).

<http://dx.doi.org/10.21523/gcj3.18020202>

© 2018 GATHA COGNITION® All rights reserved.

Analyzed data of groundwater can be used to find out its suitability for drinking, agriculture and industrial purposes (Babiker et al., 2007). This can be analyzed easily with the help of geospatial techniques in GIS [Geographic Information System] (Mahalingam et al., 2014). GIS is useful tool to analysis the changes in land use/land cover, lineament and groundwater qualities and for monitoring the groundwater resources (Chaudhary et al., 1996; Skidmore et al., 1997; Saraf, 1999; Epstein et al., 2002; Tjandra, et al., 2003; Shankar et al., 2010; Aravindan and Shankar, 2011; Wagh et al., 2016; Kadam et al., 2018).

Gaikwad et al. (2012b) have analyzed physicochemical elements in groundwater from downstream part of the Vel River Basin and reported necessity of more attention due to more industrialization, growing residential area, having water scarcity problem with zones of hard to very hard water with many kidney stone patients. Therefore, this study was undertaken with spatial analysis of groundwater qualities using GIS and RS techniques in and around Shikrapur and Talegaon Dhamdhare, Pune district, Maharashtra (India). The study is focused on analysis of water related issues.

2 STUDY AREA

Study area is downstream part of the Vel River basin (9 Km²), a tributary of Bhima River flowing through Rajgurunagar and Shirur tehsils of Pune district (Figure 1)

(Shikrapur and Talegaon Dhamdhare). The extents are from 74°07'37.686" to 74°09'47.506"E and 18°39'43.04" to 18°41'53.555"N (SOI toposheets- 47 J/2, scale- 1:50,000).

Annual rainfall is about 650 mm, receives from June to October (IMD, 2015). An average temperature ranges from 10 to 28°C. Gently rolling topography observed in the area and alluvial deposits mainly along the river course. Vel River (length 5866 m) is the major river flowing with two tributaries of 2665m and 1100m lengths (Figure 1).

The slope is estimated from ASTER images and varies 15% to 55% (Figure 2). Gentle slopes result in slower runoff and less rate of erosion with good recharge potentials (Kadam et al., 2012; Mundalik et al., 2018).

Land use and land cover (LULC) is the key factor of availability and distribution of groundwater with quality (Ghorbani et al., 2017; Singhai et al., 2017; Kadam et al., 2017). It is helpful for identifications of infiltration zones and quantification of surface runoff (Jasrotia et al., 2006; Dinesh Kumar et al., 2007). LULC modifies evapotranspiration patterns and groundwater recharge (Jasrotia et al., 2006; Manap et al., 2013).

LULC map shows dominance of land use for agricultural activities (57%) and built-up land shows 33%, fallow land observes 7% and water bodies covers 3% of the land in the study area. This map (Figure 3) was also used as base map during collection of groundwater samples.




Figure 1. Study area: Vel River Basin




Figure 2. Slope map




Figure3. Land use and Land cover

3 GEOLOGY

The study area is a part of Central Deccan Trap, Supergroup. Indrayani Formation and Upper Ratangarh Formation are exposed (Figure 4). Indrayani Formation is typically characterized by the presence of compact, massive, porphyritic basalts where the phenocrysts are embedded in fine-grained groundmass (GSI, 2001). Jointing nature have attributed them to be moderately good aquifers (Figure 4). They are seen occurring in low-lying flat plain areas of Vel River basin (GSI, 2001). Upper Ratangarh Formation is consisting of 3 compound pahoe-hoe flows (50-220m) thick in the area under investigation. It comprises mainly dark grey, fine to medium grained, massive hard and compact, non-porphyritic to porphyritic in nature with olivine phenocrysts (Godbole *et al.*, 1996; GSI, 2001).

4 METHODOLOGY

Sampling and Chemical Analysis

Sites for groundwater sampling were decided with consideration of geological and geomorphological variation from the study area (Figure 6). Thirty (30) groundwater samples from dug wells and bore wells, tapping diverse aquifers were collected in the month of April-2015 for major physicochemical analysis. The standard methods were adapted for collection and analyses of water samples given by APHA (1998). Anions and cations in groundwater samples were analyzed on High Performance Ion Chromatography (HPIC) and Inductively Coupled Plasma Atomic Emission Spectroscopy (ICPAES-9000; Shimadzu,

Made in Japan) respectively, available in Geology department of Savitribai Phule Pune University, Pune (India).




Figure 4. Geology (GSI, 2001)




Figure 5. Methodology




Figure 6. Sampling Locations

5 RESULTS AND DISCUSSIONS

5.1 Distribution of pH and EC

Groundwater in the region is slightly alkaline in nature (range: pH 7.12-8.19; average: 7.12) (Table 1) (Figure 7). The Electrical Conductivity (EC) is high ($>1900 \mu\text{S}/\text{cm}$) reported in the Central part, while lower values are estimated in the Northwest part of the region (Figure 8). Average value of EC ($781 \mu\text{mhos}/\text{cm}$) shows high mineralization in the groundwater (Pawar et al., 2008; Rubasinghe et al., 2015).

5.2 Distribution of Cations

Increase in concentration of Sodium (Na) (Figure 9) from Northwest to Southeast parts due to changes in hydro-geomorphological settings (e.g., slope) (Pawar et al., 2008) and groundwater flow except in Central part where highest Na found due to more rock-water interaction (Hem, 1959).

In many samples, low concentration of Potassium (K) is observed due to non-appearance of Potassium rich minerals (Subbarao et al., 1994; Gaikwad et al., 2019). However, increase of K in groundwater in downstream part of the study area (Figure 10) is due to use of high fertilizers and anthropogenic activities (Pawar et al., 2008; Kale et al., 2010).

Mafic minerals are principal components in the lithology from the study area contributing towards more Magnesium (Mg) as one of the major dissolved constituent in the groundwater (Hem, 1959; Subramani et al., 2010). Concentration of Mg in groundwater from study area varies from 22.42 to 198.9mg/l (Figure 11). A closely spaced contour occurs in the lower part of study area implies high salinity zone. Increase in Ca (Figure 12) in the South-Central part where deeper rock-water nitration could be the source for Ca in groundwater (Hem, 1959). Thus, Na+Ca representing 61.37% of total cations denoting major supply from weathering of plagioclase feldspar (Pawar et al., 2008). On the other hand, Ca+Mg values account 67.92% of the total cations contributing from weathering of olivine and pyroxene (Pawar et al., 2008).

5.3 Distribution of Anions

Chloride is the major anion (range: 1.43 to 781.76 mg/l; average: 335.40 mg/l) in groundwater of the study area. Higher values of chloride are reported in the Central and Southern parts (Figure 13). Cl concentration increases from upstream and downstream region (Pawar et al., 2008). Higher concentrations of chloride (Figure 13) found i. e. 781.7, 756.7 and 737.51mg/l in BW-4, BW-3 and DW-9, respectively, may be due to the accumulation of industrial waste in downstream part (Hem, 1959).

Higher values of HCO_3 are observed in South-Central part of study area while lesser values noted towards Northern side (Figure 14). More HCO_3 in groundwater is contributed from silicate and carbonate

chemical weathering (Pawar et al., 2008; Chidambaram et al., 2011). Sulphate concentrations are predominantly higher in downstream region while in upstream region shows very less recharge of groundwater (Figure 15). Anomalous higher values of Sulphate in Central part pointing towards higher fertilizers used for irrigated agriculture (Pawar et al., 2008).

The concentration of nitrate in groundwater shows range from 3.20 to 587.71mg/l and very high compare to the permissible standards by WHO. The source of NO_3 in groundwater is solid and liquid waste from human and livestock activities including agricultural runoff (Pawar et al., 2008; Brindha et al., 2010; Brindha and Elango, 2014).

Total hardness is ranges from 204.94 to 1495.47mg/l (Figure 17). Hardness exceeds total Alkalinity, it expresses the presence of both Calcium hardness and noncarbonated hardness (Chow, 1964; Subbarao et al., 2005).

5.4 Quality of Groundwater for Drinking Purposes

Water used for drinking and household purposes should not have toxic chemicals and pathogens (Nag and Das, 2017). World Health Organization (WHO, 1997) has given the standard limits for quality of water used for drinking purpose. The groundwater data compared with the WHO (1997) standards pinpointing that, except Electrical Conductivity (EC), Calcium (Ca), Potassium (K), Magnesium (Mg), Chloride (Cl) and Nitrate (NO_3) all elements are showing low to optimum values (Table 2).

Table1. Physicochemical analysis

Parameters	Maximum	Minimum	Average	Variance	Standard Deviation
pH	8.19	7.12	7.65	0.07	0.26
EC	1921.00	32.00	781.80	460992.79	678.96
Hardness	1495.47	204.94	583.19	172159.79	326.97
Na	245.40	20.97	120.43	3916.96	62.58
K	87.57	3.84	14.57	489.66	22.12
Ca	270.90	45.10	105.08	3250.09	57.00
Mg	198.90	22.42	77.51	2096.97	45.79
Cl	781.76	1.43	335.40	38415.9	195.99
HCO_3	510.00	205.00	313.50	4922.67	70.16
SO_4	331.09	20.05	100.01	8488.90	92.13
NO_3	587.71	3.20	104.64	25198.21	158.73

All values given are in mg/l, except EC in $\mu\text{mhos/cm}$ and pH on scale.

Table 2. Drinking water quality analysis (WHO, 1997)

Parameter	Permissible limit	Samples above permissible limit (%)
pH	6.5-8.5	0
EC	1500 $\mu\text{mhos/cm}$	27
Hardness	500-1500mg/l	0
Na	200-600 mg/l	0
K	30mg/l	13
Ca	75-200 mg/l	10
Mg	50-150 mg/l	13
Cl	200-600 mg/l	17
SO ₄	200-400 mg/l	0
HCO ₃	-----	----
NO ₃	50 mg/l	40




Figure 7. Distribution of pH




Figure 8 Distribution of Electrical Conductivity (EC)




Figure 9. Distribution of Sodium (Na)




Figure10. Distribution of Potassium (K)




Figure 11. Distribution of Calcium (Ca)








Figure 12. Distribution of Magnesium (Mg)

Figure 13. Distribution of Chloride (Cl)Figure 14. Distribution of Alkalinity (HCO_3)Figure 15. Distribution of Sulphate (SO_4)Figure 16. Distribution of Nitrate (NO_3)




Figure 17. Distribution of Total Hardness (TH)

6 CONCLUSION

The major ion geochemistry of groundwater from Shikrapur and Talegaon Dhamdhare area were studied for pre-monsoon season (2015).

1. Abundance of the major ions in groundwater: $\text{Na} > \text{Ca} > \text{Mg} > \text{K}$ and $\text{Cl} > \text{HCO}_3 > \text{NO}_3 > \text{SO}_4$.
2. The pH of groundwater is slightly alkaline while higher EC pinpointing to higher mineralization of groundwater.
3. Higher Sodium in groundwater from Central part anomalously due to more rock-water interaction.
4. The source of Ca and Mg in groundwater is from lithology and intensive agricultural activities in the region.
5. Weathering of light coloured minerals like plagioclase feldspar gives Sodium and Calcium (61.37 %) while Ca+Mg values accounts 67.92% of total cations supplied from weathering of dark coloured minerals like olivine and pyroxene.
6. The sources of Sulphate (SO_4) and Nitrate (NO_3) are anthropogenic and their concentration is increasing in downstream part of the study area.
7. The results show higher level of EC, Ca, Mg, K, Cl and NO_3 compared to WHO norms. Other elements show low to optimum level and good quality for drinking purposes excluding some pockets in downstream area.
8. Regular quality monitoring of groundwater with addition of rainwater harvesting is suggested to improve the quality of groundwater.

9. In nutshell, the groundwater quality from Shikrapur and Talegaon Dhamdhare areas is controlled by natural as well as anthropogenic activities.

10. The findings of the study will be helpful for monitoring the groundwater qualities in the region.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

Authors acknowledge, Head, Geology Department of Savitribai Phule Pune University (SPPU) for providing the laboratorial facility. This research project (13SCI001306: 2013-2015) was funded by BCUD-Savitribai Phule Pune University. Anonymous reviewers thanked for comments and suggestions useful to improve the manuscript.

ABBREVIATIONS

ASTER: Advanced Spaceborne Thermal Emission and Reflection Radiometer; **BW**: Bore Well; **DW**: Dug Well; **GSI**: Geological Survey of India; **IMD**: India Meteorological Department; **LULC**: Land Use Land Cover; **TH**: Total Hardness; **WHO**: World Health Organization.

REFERENCES

- Aksoy, A. O. and Scheytt, T., 2007. Assessment of groundwater pollution around Torbali, Izmir, Turkey. *Environmental geology*, 53(1), 19-25. DOI: <https://doi.org/10.1007/s00254-006-0614-5>
- APHA [American Public Health Association], 1998. Standard methods for the examination of water and wastewater. Washington DC, 20th Ed.
- Aravindan, S. and Shankar, K. J., 2011. Ground water quality maps of Paravanan River Sub Basin, Cuddalore District, Tamil Nadu, India. *Indian Soc. Remote Sens.*, 39(4), 565-581. DOI: <https://doi.org/10.1007/s12524-011-0152-9>
- Babiker, I. S., Mohamed, M. A. and Hiyama, T., 2007. Assessing groundwater quality using GIS. *Water Resources Management*, 21(4), 699-715. DOI: <https://doi.org/10.1007/s11269-006-9059-6>
- Brindha, K., and Elango, L., 2014. Soil and groundwater quality with reference to nitrate in a semiarid agricultural region. *Arabian Journal of Geosciences*, 7(11), 4683-4695. DOI: <https://doi.org/10.1007/s12517-013-1100-5>
- Brindha, K., Rajesh, R., Murugan, R. and Elango, L., 2010. Natural and anthropogenic influence on the fluoride and nitrate concentration of groundwater in parts of Nalgonda district, Andhra Pradesh, India. *Journal of Applied Geochemists*, 12(2), 231-241.
- Chaudhary, B. S., Kumar, M., Roy, A. K., and Ruhel, D. S., 1996. Applications of RS and GIS in groundwater investigations in Sohna block, Gurgaon district, Haryana, India. *International Archives of Photogrammetry and Remote Sensing*, 31, 18-23.
- Chidambaram, S., Karmegam, U., Prasanna, M. V., Sasidhar, P., and Vasanthavigar, M. 2011. A study on hydrochemical elucidation of coastal groundwater in and around Kalpakkam region, Southern India.

- Environmental Earth Sciences*, 64(5), 1419-1431. DOI: <https://doi.org/10.1007/s12665-011-0966-3>
- Chow, V. T., 1964. Hydrology and its development. *Handbook of Applied Hydrology*, 1-1.
- Dinesh Kumar, P. K., Gopinath, G. and Seralathan, P. 2007. Application of remote sensing and GIS for the demarcation of groundwater potential zones of a river basin in Kerala, southwest coast of India. *International Journal of Remote Sensing*, 28(24), 5583-5601. DOI: <https://doi.org/10.1080/01431160601086050>
- Epstein, J., Payne, K. and Kramer, E., 2002. Techniques for mapping suburban sprawl. *Photogrammetric engineering and remote sensing*, 68(9), 913-918.
- Gaikwad, S. K. and Pawar, N. J., 2012a. Imprints of Lithological Diversity on the Chemical Composition of Groundwater from Sindhudurg District, Maharashtra. *Mem Geological Society of India*, 60, 109-126.
- Gaikwad, S. K., Kouhsari, M. and Pawar, N. J., 2012b. Geochemistry of groundwaters from the Upper Vel River Basin, part of Pune District, Maharashtra. *Fifth International Groundwater Conference (IGWC-2012)*, December 18-21, Aurangabad, India.
- Gaikwad, S., Gaikwad, S., Meshram, D., Wagh, V., Kandekar, A. and Kadam, A., 2019. Geochemical mobility of ions in groundwater from the tropical western coast of Maharashtra, India: implication to groundwater quality. *Environment, Development and Sustainability*, 1-34. DOI: <https://doi.org/10.1007/s10668-019-00312-9>
- Ganyaglo, S. Y., Banoeng-Yakubo, B., Osa, S., Dampare, S. B. and Fianko, J. R., 2011. Water quality assessment of groundwater in some rock types in parts of the eastern region of Ghana. *Environmental Earth Sciences*, 62(5), 1055-1069. DOI: <https://doi.org/10.1007/s12665-010-0594-3>
- Ghorbani N. S., Falah, F., Daneshfar, M., Haghizadeh, A. and Rahmati, O., 2017. Delineation of groundwater potential zones using remote sensing and GIS-based data-driven models. *Geocarto international*, 32(2), 167-187. DOI: <https://doi.org/10.1080/10106049.2015.1132481>
- Godbole, S. M., Rana, R. S. and Natsu, S. R., 1996. Lava stratigraphy of Deccan basalts of Western Maharashtra. *Gondwana Geol. Mag. Spl.* 2, 125-135.
- GSI [Geological Survey of India], 2001. District Resource Map of Pune.
- Hem, J. D., 1959. *Study and interpretation of the chemical characteristics of natural water* (No. 1473). US Government Printing Office.
- IMD [India Meteorological Department], 2015. Annual Weather Reports, Pune (India).
- Jasrotia, A. S. and Singh, R., 2006. Modeling runoff and soil erosion in a catchment area, using the GIS, in the Himalayan region, India. *Environmental Geology*, 51(1), 29-37. DOI: <https://doi.org/10.1007/s00254-006-0301-6>
- Kadam, A. K., Jaweed, T. H., Umrikar, B. N., Hussain, K. and Sankhua, R. N., 2017. Morphometric prioritization of semi-arid watershed for plant growth potential using GIS technique. *Modeling Earth Systems and Environment*, 3(4), 1663-1673. DOI: <https://doi.org/10.1007/s40808-017-0386-9>
- Kadam, A. K., Kale, S. S., Pande, N. N., Pawar, N. J. and Sankhua, R. N., 2012. Identifying potential rainwater harvesting sites of a semi-arid, basaltic region of Western India, using SCS-CN method. *Water resources management*, 26(9), 2537-2554. DOI: <https://doi.org/10.1007/s11269-012-0031-3>
- Kadam, A., Karnewar, A. S., Umrikar, B. and Sankhua, R. N., 2018. Hydrological response-based watershed prioritization in semiarid, basaltic region of western India using frequency ratio, fuzzy logic and AHP method. *Environment, Development and Sustainability*, 1-25. DOI: <https://doi.org/10.1007/s10668-018-0104-4>
- Kale, S. S., Kadam, A. K., Kumar, S. and Pawar, N. J., 2010. Evaluating pollution potential of leachate from landfill site, from the Pune metropolitan city and its impact on shallow basaltic aquifers. *Environmental monitoring and assessment*, 162(1-4), 327-346. DOI: <https://doi.org/10.1007/s10661-009-0799-7>
- MacDonald, A. M. and Davies, J., 2000. A brief review of groundwater for rural water supply in sub-Saharan Africa.
- Mahalingam, B., Bhauso, R., Deepali, M. and Jayashree, P., 2014. Assessment of Groundwater Quality Using GIS Techniques: A Case Study of Mysore City. *International Journal of Engineering and Innovative Technology (IJEIT)*, 117-122.
- Manap, M. A., Sulaiman, W. N. A., Ramli, M. F., Pradhan, B. and Surip, N., 2013. A knowledge-driven GIS modeling technique for groundwater potential mapping at the Upper Langat Basin, Malaysia. *Arabian Journal of Geosciences*, 6(5), 1621-1637. DOI: <https://doi.org/10.1007/s12517-011-0469-2>
- Mundalik, V., Fernandes, C., Kadam, A. K. and Umrikar, B. N., 2018. Integrated geomorphological, geospatial and AHP technique for groundwater prospects mapping in Basaltic Terrain. *Hydrospatial Analysis*, 2, 16-27. DOI: <http://dx.doi.org/10.21523/gcj3.18020102>
- Nag, S. K. and Das, S., 2017. Assessment of groundwater quality from Bankura I and II Blocks, Bankura District, West Bengal, India. *Applied Water Science*, 7(6), 2787-2802. DOI: <https://doi.org/10.1007/s13201-017-0530-8>
- Nathanson, J., A., 1986. *Basic Environmental Technology: Water Supply, Waste Disposal and Pollution Control*. John Wiley.
- Pawar, N. J., Pawar, J. B., Kumar, S. and Supekar, A., 2008. Geochemical eccentricity of ground water allied to weathering of basalts from the Deccan Volcanic Province, India: insinuation on CO₂ consumption. *Aquatic Geochemistry*, 14(1), 41-71. DOI: <https://doi.org/10.1007/s10498-007-9025-9>
- Ravikumar, P. and Somashekar, R. K., 2013. A geochemical assessment of coastal groundwater quality in the Varahi river basin, Udupi District, Karnataka State, India. *Arabian Journal of Geosciences*, 6(6), 855-1870. DOI: <https://doi.org/10.1007/s12517-011-0470-9>
- Ravikumar, P., Venkatesharaju, K. and Somashekar, R. K., 2010. Major ion chemistry and hydrochemical studies of groundwater of Bangalore South Taluk, India. *Environmental monitoring and assessment*, 163(1-4), 643-653. DOI: <https://doi.org/10.1007/s10661-009-0865-1>
- Rubasinghe, R., Gunatilake, S. K. and Chandrajith, R., 2015. Geochemical characteristics of groundwater in different climatic zones of Sri Lanka. *Environmental earth sciences*, 74(4), 3067-3076. DOI: <https://doi.org/10.1007/s12665-015-4339-1>
- Saraf, A. K., 1999. *A report on Land use Modelling in GIS for Bankura District. DST, NRDMS*.
- Shankar, K., Aravindan, S. and Rajendran, S., 2010. GIS based groundwater quality mapping in Paravanan river sub-basin, Tamil Nadu, India. *International Journal of Geomatics and Geosciences*, 1(3), 282.
- Singhai, A., Das, S., Kadam, A. K., Shukla, J. P., Bundela, D. S. and Kalashetty, M., 2017. GIS-based multi-criteria approach for identification of rainwater harvesting zones

- in upper Betwa sub-basin of Madhya Pradesh, India. *Environment, Development and Sustainability*, 1-21. DOI: <https://doi.org/10.1007/s10668-017-0060-4>
- Skidmore, A. K., Bijker, W., Schmidt, K. and Kumar, L., 1997. Use of remote sensing and GIS for sustainable land management. *ITC journal*, 3(4), 302-315.
- Subbarao, K. V., 1994. Stratigraphy and structure of parts of the central Deccan basalt province: eruptive models. *Volcanism*, 321-332.
- Subbarao, N. S., Nirmala, I. S. and Suryanarayana, K., 2005. Groundwater quality in a coastal area: a case study from Andhra Pradesh, India. *Environmental Geology*, 48(4-5), 543-550. DOI: <https://doi.org/10.1007/s00254-005-1306-2>
- Subramani, T., Rajmohan, N. and Elango, L., 2010. Groundwater geochemistry and identification of hydrogeochemical processes in a hard rock region, Southern India. *Environmental monitoring and assessment*, 162(1-4), 123-137. DOI: <https://doi.org/10.1007/s10661-009-0781-4>
- Tjandra, F. L., Kondhoh, A. and Mohammed, M. A., 2003. A conceptual database design for hydrology using GIS. *Proceedings of Asia Pacific association of hydrology and water resources, Kyoto*, 13-15.
- Wagh, V. M., Panaskar, D. B., Mukate, S. V., Gaikwad, S. K., Muley, A. A. and Varade, A. M., 2018b. Health risk assessment of heavy metal contamination in groundwater of Kadava River Basin, Nashik, India. *Modeling Earth Systems and Environment*, 4(3), 969-980. DOI: <https://doi.org/10.1007/s40808-018-0496-z>
- Wagh, V. M., Panaskar, D. B., Varade, A. M., Mukate, S. V., Gaikwad, S. K., Pawar, R. S. Muley A. A. and Aamalawar, M. L., 2016. Major ion chemistry and quality assessment of the groundwater resources of Nanded tehsil, a part of southeast Deccan Volcanic Province, Maharashtra, India. *Environmental Earth Sciences*, 75(21), 1418. DOI: <https://doi.org/10.1007/s12665-016-6212-2>
- Wagh, V., Panaskar, D., Muley, A., Mukate, S. and Gaikwad, S. 2018a., Neural network modelling for nitrate concentration in groundwater of Kadava River basin, Nashik, Maharashtra, India. *Groundwater for Sustainable Development*, 7, 436-445. DOI: <https://doi.org/10.1016/j.gsd.2017.12.012>
- WHO [World Health Organization], 1997. *Guidelines for Drinking Water Quality Health Criteria and other Supporting Information*. 2, 2nd Ed., WHO, Geneva.
